

Web Images Videos Maps News Shopping Gmail more ▾

Sign in

Google scholar [Advanced Scholar Search](#)
[Scholar Preferences](#)

Scholar ☐ - Results **1 - 20** of about **68**

[PDF] Utilization and predictability in scheduling the IBM SP2 with backfilling

DG Feitelson, AM Weil - 12th International Parallel Processing ..., 1998 - Citeseer

... the jobs in the queue, that is, to use non-FCFS policies [4]. Consider the following scenario, where a number of jobs are **running side by side**, and the ... Using this information, only short jobs — that are expected to **terminate** in time — are allowed to leapfrog a waiting large job. ...

Cited by 170 - Related articles - View as HTML - All 9 versions

[CITATION] The effects of **termination** of the CS and avoidance of the US on avoidance learning.

LJ Kamin - Journal of Comparative and Physiological Psychology, 1956

Cited by 54 - Related articles - All 6 versions

A rapid method for determining sequences in DNA by primed synthesis with DNA polymerase* 1

F Sanger, AR Coulson - Journal of Molecular Biology, 1975 - Elsevier

... Incubation was at 0C. After 2 min 50 id were removed and added to 5 il 02 MEDTA to **terminate** the reaction. After ... radioautograph. This is easy where samples are **run side by side** but involves careful measuring for ones further apart. ...

Cited by 920 - Related articles - All 3 versions

[CITATION] Utilization, predictability, workloads, and user runtime estimates in scheduling the IBM SP2 with backfilling

AW Mu'aleem, DG Feitelson - IEEE Transactions on Parallel and Distributed Systems, 2001

Cited by 310 - Related articles - BL Direct - All 14 versions

[CITATION] Does CS **termination** reinforce avoidance behavior

RC Bolles, LW Stokes, MS ... - Journal of ..., 1966 - American Psychological Association ...

Cited by 63 - Related articles - All 5 versions

The **termination** of ascending tracts in the thalamus of the macaque monkey

WE Le Gros Clark - Journal of Anatomy, 1936 -.ncbi.nlm.nih.gov

... 2 A). The lesion thus interrupts the fibres which **run** medially from the ventro-lateral half of the lower part of the nucleus. ... Left **side**. As the sections are traced rostrally, the lesion on this **side** is first seen entering the **termination** of the fasciculus cuneatus at the level of the lower ...

Cited by 100 - Related articles - All 2 versions

Observation on the supra-optic decussations, in the albino rat

JM Minderhoud - The Journal of Comparative Neurology, 1967 - interscience.wiley.com

... medial **side** and ascended to the supra-optic decussa- tion on the dorso-caudal **side** of the ... in the medulla oblongata were not followed by degeneration of any additional fibers **running** to the ... with another localization in the whole fiber tract or of the place of **termination**, then the ...

Cited by 23 - Related articles - All 2 versions

Modular **termination** system for communication lines

WA Brey - US Patent 4,160,880, 1979 - Google Patents

... distribution terminal 36. Accordingly, as seen in FIG. 1, from each other for locating the cables **side-by-side** as modular **termination** systems 24 according to the inven- they **run** in a direction away from the field. The cable 30 tion ...

[Cited by 23](#) - [Related articles](#)

Termination of transcription in vitro in the escherichia coli tryptophan operon leader region* 1

F Lee, CL Squires, C Squires, C Yanofsky - Journal of Molecular Biology, 1976 - Elsevier

... In addition, samples treated with alkaline phosphatase were **run side by side** on DEAEpaper (Whatman DE81, pH 35) with untreated samples, along with a series of oligonucleotides of known length with the general sequence (Up)Gp (IGpUsGp). **TERMINATION** OF trp ...

[Cited by 65](#) - [Related articles](#) - [All 2 versions](#)

[PDF] The pharynx of Caenorhabditis elegans

DG Albertson, JN Thomson - ... Transactions of the Royal Society of London. ..., 1976 - JSTOR

... **run** anteriorly in the subventral nerve cords forming synapses on muscles m5 and pass directly through the nerve ring to the most anterior projection of muscles m5. These then travel dorsally forming motor synapses on muscles m4. They meet on the dorsal **side** and **terminate** in ...

[Cited by 353](#) - [Related articles](#) - [All 4 versions](#)

[PDF] The local extension of nerve fibres into denervated areas of skin

G Weddell, L Gutfmann, E Gutfmann - Journal of Neurology and ..., 1941 - jnnp.bmj.com

... less marked still, though in all cases some slight difference between the two sides could be detected in favour of the unoperated **side**. ... There is centrally placed a proximo-distally **running** nerve trunk, from which spring subsidiary nerve trunks, giving rise to bundles which scatter ...

[Cited by 146](#) - [Related articles](#) - [All 7 versions](#)

[CITATION] Continual queries for internet scale event-driven informationdelivery

L Liu, C Pu, W Tang - IEEE Transactions on Knowledge and Data ..., 1999

[Cited by 365](#) - [Related articles](#) - [BL Direct](#) - [All 29 versions](#)

Muscle receptor organs in the abdomen of Homarus vulgaris and Palinurus vulgaris

JS Alexandrowicz - Journal of Cell Science, 1951 - jcs.biologists.org

... 178 Ahxandrovncz—Muscle Receptor Organs in the the surrounding connective tissue fibres but mainly **run** longitudinally, vidently having the function of intercalated tendons. The muscles coming from each **side terminate** in them somewhat unevenly, so that serial sections ...

[Cited by 202](#) - [Related articles](#) - [All 2 versions](#)

Antennal movements induced by odour and central projection of the antennal neurones in the honey-bee

H Suzuki - Journal of Insect Physiology, 1975 - Elsevier

... Injection of the fluorescent dye into the antennal nerve after degeneration of the antennal sensory neurones showed that the antennal motoneurones **run** in the ventral **side** of the antennal and dorsal lobes, and **terminate** in the marginal region of the ipsilateral oesophageal ...

[Cited by 78](#) - [Related articles](#)

Gang scheduling performance benefits for fine-grain synchronization

DG Feltelson, L Rudolph - Journal of Parallel and Distributed Computing, 1992 - Elsevier

... As a **side** note, we observe that twophase blocking can be used to place a bound on ... The system may be partitioned into two main layers: A **run** time library that supports the ParC ... Thus they **terminate** early in the measurement, and serve to cause the different processors to start ...

[Cited by 232](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)

Rules for alpha-helix termination by glycine

R Aurora, R Srinivasan, GD Rose - Science, 1994 - sciencemag.org

... than a computer algorithm can be deducedby the observa- tion of the way a **running** program trans ... 5), the NH2-termini of helices were capped predominantly by nearby residue **side** chains ... Approximately one-third ofall a helices **terminate** with a Gly residue at the COOH- terminus ...

[Cited by 196](#) - [Related articles](#) - [BL Direct](#) - [All 9 versions](#)

Organization of the stomatogastric ganglion of the spiny lobster

DM Maynard, AI Selverston - Journal of Comparative Physiology A: ..., 1975 - Springer
 ... Physiol., Vol. 100 Page 6. 166 DM Maynard and AI Selverston containing the VD and the other
 the IC but in most cases the two axons **run side-by-side** down a single nerve. ... Its axon is found
 in the stomatogastric nerve (SGN) and its final **termination** is unknown. ...

[Cited by 80](#) - [Related articles](#)

An operational semantics for StAC, a language for modelling long-running business transactions

M Butler, C Ferreira - Coordination Models and Languages, 2004 - Springer
 ... In the case that a compensation pair is **running** in parallel with a process that executes an early
termination, this early **termination** cannot ... of a process N (where $N = P$ is an equation) will substitute
 call(N) by the process on the left-**side** of the ... to skip) when both P and Q **terminate**: ...

[Cited by 107](#) - [Related articles](#) - [BL Direct](#) - [All 14 versions](#)

An experimental study of the course and **termination** of the projection fibers from cortical areas 4 and 6 in the cat


K Niimi, S Kishi, M Miki, S Fujita - Psychiatry and Clinical ..., 1963 - interscience.wiley.com
 ... has been no precise description as to whether fibers from the motor cortex **terminate** in the ... On
 the **side** contralateral to the lesion, degenerating fibers are not observed in the centre ... In case
 11, a few fine beaded degenerating axons **running** dorsomedially and dorsally are found ...

[Cited by 62](#) - [Related articles](#) - [All 3 versions](#)

Backfilling with lookahead to optimize the performance of parallel job scheduling

E Shmueli, D Feitelson - Job Scheduling Strategies for Parallel ..., 2003 - Springer
 ... started. Waiting jobs which are expected to **terminate** after the shadow time can use
 these extra processors, referred to as the shadow free capacity, and **run side by**
side together with wj1, without effecting its start time. As long ...

[Cited by 62](#) - [Related articles](#) - [BL Direct](#) - [All 8 versions](#)

 [Create email alert](#)

Google 

Result Page: [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [Next](#)

[Go to Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2010 Google